

Improving astrophysical γ -ray modeling with Gaussian processes and variational inference

Siddharth Mishra-Sharma and Kyle Cranmer, New York University




NYU

Center for Cosmology and Particle Physics

Motivation

- Signals of dark matter (DM) may be hiding in γ -ray observations of the Galactic Center
- Galactic backgrounds of astrophysical origin make up a large fraction ($\gtrsim 80\%$) of the data in this region
- The poorly-understood spatial morphology of these backgrounds makes it difficult to characterize DM signals [1-2]
- Motivates analysis techniques that can account for uncertainty in knowledge of Galactic background contribution (e.g., Refs. [3,5])




Summary

- Traditional γ -ray analyses model data as a sum of rigid spatial templates, each corresponding to a specific astrophysical component [4]
- We modulate the poorly-understood Galactic background templates by a **Gaussian process (GP)** in order to give them more freedom and account for uncertainty in their spatial variation
- The GP is included as part of a larger probabilistic model that includes parameters describing other modeled components, including a dark matter signal. Variational inference is used for tractable analysis.

Traditional template fitting


- Spatially-binned (pixelized) data d^p is modeled as a Poisson realization of sum of templates T_i^p
- Spatial profile of each template is rigidly fixed
- Normalizations A_i of templates are floated as free parameters of the model

$$d^p \sim \text{Pois} \left(\sum_i A_i T_i^p \right)$$





(poorly-understood) Augmenting Galactic background template with a GP

- Modulate Galactic background template with GP
- Exponential link function to ensure positivity





Proof-of-principle analysis on simulated data

- Create simulated data using one Galactic background model (Model 1), analyze using other model (Model 2) (templates from Refs. [5-6])
- GP used for variational posterior defined using GPyTorch [7] and Pyro [8]
- Variational distributions of non-GP parameters (template normalizations) $q(A_i | \text{GP})$ defined using inverse autoregressive flows conditioned on GP summary statistics in order to capture correlations between GP and non-GP parameters



Ratio of Galactic background templates Model 1 (in simulation) and Model 2 (in analysis), inferred via GP (—) vs truth (—)



Machine Learning and the Physical Sciences Workshop 2020



References

- | | |
|----------------------------------|--------------------------------------|
| [1] Leane & Slatyer [1904.08430] | [5] Buschmann et al [2002.12373] |
| [2] Leane & Slatyer [2002.12370] | [6] Mishra-Sharma et al [1612.03173] |
| [3] Storm et al [1705.04065] | [7] Gardner et al [1809.11165] |
| [4] Lee et al [1506.05124] | [8] Bingham et al [1810.09538] |

Siddharth Mishra-Sharma and Kyle Cranmer
Semi-parametric γ -ray modeling with Gaussian processes and variational inference
[2010.10450]

<https://github.com/smsharma/gamma-gp>

